



Carson



Carson is a high yielding wax specialty bean with improved characteristics and disease package. Producing a slimmer shape than Indy Gold, in trials Carson has delivered a high percentage of refined 4 sieve pods approximately two days earlier. Carson adds to these strengths with an excellent disease package including bean common mosaic virus, Anthracnose and resistance to bacterial brown spot.

T R I A L D A T A *	
PVP Status	A
Approx. Days to Maturity	56
Heat Units (C°/F°)	638/1148
Approx. Length(in./cm.)	5.3/13.5
Average Sieve Size	3.8
Pod Color	Yellow
Disease Resistance	HR: BCMV, CI IR: BCTV, Pss

* See Back Side for Disease Resistance Descriptions



Note: All variety information presented herein is based on field and laboratory observation. Actual crop yield and quality are dependent upon many factors beyond our control and NO WARRANTY is made for crop yield and quality. Since environmental conditions and local practices may affect variety characteristics and performance, we disclaim any legal responsibility for these. Read all tags and labels. They contain important conditions of sale, including limitations of warranties and remedies. ROGERS® is a registered trademark of a Syngenta Group Company. Syngenta Seeds, Inc., P.O. Box 4188, Boise, ID 83711-4188, U.S.A. www.rogersadvantage.com



Trial Data

KEY TO RESISTANCE ABBREVIATIONS FOR BEAN	
BCMV	Bean common mosaic caused by the specified strains of <i>Bean common mosaic virus</i>
BCTV	Curly top caused by <i>Beet curly top virus</i>
BGYMV	Bean golden yellow mosaic caused by <i>Bean golden yellow mosaic virus</i>
Cl	Anthrachnose caused by <i>Colletotrichum lindemuthianum</i>
Psp	Halo blight caused by <i>Pseudomonas savastanoi</i> pv. <i>phaseolicola</i>
Pss	Bacterial brown spot caused <i>Pseudomonas syringae</i> pv. <i>syringae</i>
Ua	Rust caused by the specified races of <i>Uromyces appendiculatus</i>
HR	High Resistance: describes plant varieties that highly restrict the growth and development of the specified pest or pathogen under normal pest or pathogen pressure when compared to susceptible varieties. Highly resistant varieties may, however, exhibit some symptoms or damage under heavy pest or pathogen pressure.
IR	Intermediate Resistance: describes plant varieties that restrict the growth and development of the specified pest or pathogen, but may exhibit a greater range of symptoms or damage compared to highly resistant varieties. Intermediately resistant varieties will still show less severe symptoms or damage than susceptible plant varieties when grown under similar environmental conditions and/or pest or pathogen pressure.
In cases where specific races or strains are not noted the variety is resistant to some, but not necessarily all known races or strains of the pathogen.	
Note: All variety information presented herein is based on field and laboratory observation. Actual crop yield, quality, and level of claimed pest and pathogen resistances, are dependent upon many factors beyond our control and NO WARRANTY is made for crop yield, quality, and level of claimed pest and pathogen resistances. Since environmental conditions and local practices may affect variety characteristics and performance, we disclaim any legal responsibility for these. Read all tags and labels. They contain important conditions of sale, including limitations of warranties and remedies. Making Superior Vegetables a Reality™ is a trademark of a Syngenta Group Company. ROGERS® is a registered trademark of a Syngenta Group Company.	