



# Renegade



A new standard for dark beans hits the stage with Renegade. With Renegade you can meet customers expectation for extremely dark color pods. Designed for the Western U.S. and export market in Mexico, Renegade has an upright plant that may be machine or hand harvested.

T R I A L   D A T A *	
<b>PVP Status</b>	A
<b>Approx. Days to Maturity</b>	56
<b>Heat Units (C°/F°)</b>	628/1131
<b>Approx. Length(in./cm.)</b>	5.5/14
<b>Average Sieve Size</b>	4.0
<b>Pod Color</b>	Very dark green
<b>Disease Resistance</b>	<b>HR:</b> BCMV <b>IR:</b> BCTV

\* See Back Side for Disease Resistance Descriptions



Renegade



Note: All variety information presented herein is based on field and laboratory observation. Actual crop yield and quality are dependent upon many factors beyond our control and NO WARRANTY is made for crop yield and quality. Since environmental conditions and local practices may affect variety characteristics and performance, we disclaim any legal responsibility for these. Read all tags and labels. They contain important conditions of sale, including limitations of warranties and remedies. ROGERS® is a registered trademark of a Syngenta Group Company. Syngenta Seeds, Inc., P.O. Box 4188, Boise, ID 83711-4188, U.S.A. [www.rogersadvantage.com](http://www.rogersadvantage.com)



## Trial Data

KEY TO RESISTANCE ABBREVIATIONS FOR BEAN	
<b>BCMV</b>	Bean common mosaic caused by the specified strains of <i>Bean common mosaic virus</i>
<b>BCTV</b>	Curly top caused by <i>Beet curly top virus</i>
<b>BGYMV</b>	Bean golden yellow mosaic caused by <i>Bean golden yellow mosaic virus</i>
<b>CI</b>	Anthrachnose caused by <i>Colletotrichum lindemuthianum</i>
<b>Psp</b>	Halo blight caused by <i>Pseudomonas savastanoi</i> pv. <i>phaseolicola</i>
<b>Pss</b>	Bacterial brown spot caused <i>Pseudomonas syringae</i> pv. <i>syringae</i>
<b>Ua</b>	Rust caused by the specified races of <i>Uromyces appendiculatus</i>
<b>HR</b>	<b>High Resistance:</b> describes plant varieties that highly restrict the growth and development of the specified pest or pathogen under normal pest or pathogen pressure when compared to susceptible varieties. Highly resistant varieties may, however, exhibit some symptoms or damage under heavy pest or pathogen pressure.
<b>IR</b>	<b>Intermediate Resistance:</b> describes plant varieties that restrict the growth and development of the specified pest or pathogen, but may exhibit a greater range of symptoms or damage compared to highly resistant varieties. Intermediately resistant varieties will still show less severe symptoms or damage than susceptible plant varieties when grown under similar environmental conditions and/or pest or pathogen pressure.
In cases where specific races or strains are not noted the variety is resistant to some, but not necessarily all known races or strains of the pathogen.	
<b>Note:</b> All variety information presented herein is based on field and laboratory observation. Actual crop yield, quality, and level of claimed pest and pathogen resistances, are dependent upon many factors beyond our control and NO WARRANTY is made for crop yield, quality, and level of claimed pest and pathogen resistances. Since environmental conditions and local practices may affect variety characteristics and performance, we disclaim any legal responsibility for these. Read all tags and labels. They contain important conditions of sale, including limitations of warranties and remedies. Making Superior Vegetables a Reality™ is a trademark of a Syngenta Group Company. ROGERS® is a registered trademark of a Syngenta Group Company.	