



Golden Delight



Golden Delight has intermediate resistance to Zucchini yellow mosaic virus and Watermelon mosaic virus, which allows this variety to produce high quality, very uniform golden-yellow fruit with little to no greening when these viruses are present. Fruit are similar to Golden Dawn III and mature at approximately the same time. Golden Delight's vigorous, semi-spineless plant minimizes the risk of fruit damage and facilitates harvest. For a season-long quality golden zucchini program, plant Golden Dawn III early in the spring followed by Golden Delight.

T R I A L D A T A *	
Approx. Days to Maturity	47
Average Length (in.)	7 - 8
Average Diameter (in.)	1.5 - 1.75
Shape	Cylindrical
Fruit Color	Deep golden yellow
Plant Type	Upright. Open.
Disease Resistance	IR: ZYMV;WMV

* See Back Side for Disease Resistance Descriptions




Golden Delight



Note: All variety information presented herein is based on field and laboratory observation. Actual crop yield and quality are dependent upon many factors beyond our control and NO WARRANTY is made for crop yield and quality. Since environmental conditions and local practices may affect variety characteristics and performance, we disclaim any legal responsibility for these. Read all tags and labels. They contain important conditions of sale, including limitations of warranties and remedies. ROGERS® is a registered trademark of a Syngenta Group Company. Syngenta Seeds, Inc., P.O. Box 4188, Boise, ID 83711-4188, U.S.A. www.rogersadvantage.com



KEY TO RESISTANCE ABBREVIATIONS FOR SQUASH

CMV	Cucumber mosaic caused by <i>Cucumber mosaic virus</i>
Sf	Powdery mildew caused by <i>Sphaerotheca fuliginea</i>
SLCV	Squash leaf curl caused by <i>Squash leaf curl virus</i>
WMV	Watermelon mosaic caused by <i>Watermelon mosaic virus</i>
ZYMV	Zucchini yellows caused by <i>Zucchini yellow mosaic virus</i>
HR	High Resistance: describes plant varieties that highly restrict the growth and development of the specified pest or pathogen under normal pest or pathogen pressure when compared to susceptible varieties. Highly resistant varieties may, however, exhibit some symptoms or damage under heavy pest or pathogen pressure.
IR	Intermediate Resistance: describes plant varieties that restrict the growth and development of the specified pest or pathogen, but may exhibit a greater range of symptoms or damage compared to highly resistant varieties. Intermediately resistant varieties will still show less severe symptoms or damage than susceptible plant varieties when grown under similar environmental conditions and/or pest or pathogen pressure.
	The VIP seal denotes Value-added, Innovation and Performance

In cases where specific races or strains are not noted the variety is resistant to some, but not necessarily all known races or strains of the pathogen.

Note: All variety information presented herein is based on field and laboratory observation. Actual crop yield, quality, and level of claimed pest and pathogen resistances, are dependent upon many factors beyond our control and NO WARRANTY is made for crop yield, quality, and level of claimed pest and pathogen resistances. Since environmental conditions and local practices may affect variety characteristics and performance, we disclaim any legal responsibility for these. Read all tags and labels. They contain important conditions of sale, including limitations of warranties and remedies. Making Superior Vegetables a Reality™ is a trademark of Syngenta Group Company. ROGERS® is a registered trademark of Syngenta Group Company. www.rogersadvantage.com